

## **Impacto da cana de açúcar e amendoim na perda de nitrogênio na sub-bacia do rio Mogi-Guaçu**

Paulo Alexandre da Silva, Bruna Cristina de Lima Candido, Teresa Cristina Tarlé Pissarra, Alan Rodrigo Panosso, Glauco de Souza Rolim

Universidade Estadual Paulista "Júlio de Mesquita Filho" (UNESP), Jaboticabal-SP, Brasil

E-mail: paullo-alex@outlook.com

**Resumo:** Sertãozinho-SP é um grande produtor de cana-de-açúcar e amendoim mundial. Na região usa-se a técnica de rotação de culturas que tem como objetivos a recuperação dos solos e o beneficiamento comercial dos produtores. O objetivo deste estudo foi quantificar as perdas de nitrogênio, sob a influência da paisagem, seu uso e ocupação em dois sistemas de produção: amendoim e cana-de-açúcar, por meio da simulação do modelo SWAT. Foram mantidos para as duas culturas os atributos topográficos, as condições físicas e climáticas da sub-bacia hidrográfica do Rio Mogi-Guaçu, localizada no município de Sertãozinho – SP. As maiores perdas de nitrogênio (N) foram observadas na cultura do amendoim, com relação à cana (118,44 e 34,48 kg/ha, respectivamente). Para o nitrogênio orgânico, escoamento superficial de nitrato, nitrato lixiviado, fluxo lateral de nitrato e nitrato de águas subterrâneas foram registrados os valores de 0,4 e 29,9; 0,46 e 0,34; 32,74 e 85,99; 0,19 e 0,37; 0,69 e 1,84 kg/há, respectivamente. A menor quantidade de cobertura vegetal produzida pelo amendoim influenciou nas perdas de nitrogênio no solo. Os resultados mostraram que a simulação pode ser usada como uma ferramenta de gestão agrícola e para medidas políticas e ambientais, visando a conservação do solo e da água.

**Palavras-chave:** modelo SWAT; Rotação de culturas; Cana-de-açúcar e amendoim; Perdas de nitrogênio.

### **Impact of sugarcane and peanut on nitrogen losses in the Mogi-guaçu river sub-basin**

**Abstract:** Sertãozinho-SP is a major producer of sugarcane and peanuts in the world. In the region, the technique of crop rotation is used, which aims to recover soils and commercially improve producers. The objective of this study was to quantify nitrogen losses, under the influence of the landscape, its use and occupation in two production systems: peanuts and sugarcane, through the simulation of the SWAT model. The topographic attributes, physical and climatic conditions of the Mogi-Guaçu River sub-basin, located in the municipality of Sertãozinho – SP, were maintained for both cultures. The highest losses of nitrogen (N) were observed in the peanut crop, in relation to sugarcane (118.44 and 34.48 kg/ha, respectively). For organic nitrogen, nitrate runoff, nitrate leached, nitrate lateral flow and groundwater nitrate, values of 0.4 and 29.9 were recorded; 0.46 and 0.34; 32.74 and 85.99; 0.19 and 0.37; 0.69 and 1.84 kg/ha, respectively. The smaller amount of plant cover produced by peanuts influenced the loss of nitrogen in the soil. The results showed that simulation can be used as an agricultural management tool and for political and environmental measures, aiming at soil and water conservation.

**Keywords:** SWAT model; Crop rotation; Sugarcane and peanuts; Nitrogen losses.

## Introduction

Alta Mogiana region (Sertãozinho, Jaboticabal, Dumont and Ribeirão Preto) is one of the main centers for the production of peanuts and sugarcane in Brazil. In general, sugarcane producers sow peanuts in rotation. Crop rotation is a technique that provides the annual alternation of plant species in the same productive area, whose chosen crops have commercial purposes and have the capacity to recover the soil. Generally, in sugarcane reforms, rotations are carried out by species called green manures that form a surface cover and promote the maintenance of the physical, chemical and biological properties of the soil. The most used plants in the region are *Crotalaria juncea*, soybeans and peanuts. The main benefits that the crop rotation system offers are: increased productivity, control of weeds, diseases and pests, improvement in soil fertility and physical characteristics, efficiency in the use of water and nutrients, optimization of the use of machines in ownership and risk reduction of crop yield losses [1]. Besides, when plant residues are left or kept in direct contact with the soil surface, they are extremely efficient in controlling the erosion process in between furrows.

## Objectives


The aim of this study was to quantify nitrogen losses under the influence of landscape and land use and occupation of two production systems, peanuts and sugarcane, by simulating the SWAT model, keeping for both crops the topographic characteristics and the physical and climatic factors of the Mogi-Guaçu River sub-basin, located in the municipality of Sertãozinho – SP.

## Methodology

The municipality of Sertãozinho is located in the Center-North region of the State of São Paulo, located in the Planalto Ocidental Paulista, under the geographic coordinates 21° 08' 16" South Latitude and 47° 59' 25" West Longitude of Greenwich. It has a territorial area of approximately 403.89 km<sup>2</sup> [2]. According to Köppen's classification, the climate is Aw, tropical, wet summer and dry winter. The average annual temperature in the region is 22.8 °C, the average annual rainfall is around 1,588.5 mm. The predominant relief is soft-wavy and wavy, with altitude ranging from 310 m to 610 m. The Municipality is located in the Mogi-Guaçu Basin, located under the Alta Mogiana region and the predominant vegetation cover is the Cerrado [2]. The digital elevation model (DEM) of the study area was acquired through the worldwide HydroSHEDS – USGS database, whose approximate resolution is 1 kilometer.

Figure 1 shows the MDE for the municipality of Sertãozinho – SP, highlighted in red for the sub-basin under study. For the preparation of the soil map, the pedological model provided by the Brazilian Agricultural Research Corporation [3] was used, based on the Brazilian soil classification system [4]. The land use and occupation map were obtained by joining the bands provided by the Landsat-8 satellite.

The temporal resolution used was 16 days and the spatial resolution was 30 meters for the visible bands.



**Figure 01.** Digital map of digital elevation of the city of Sertãozinho – SP. Demarcation of the study sub-basin of the Mogi Guaçu basin (red). Source: Own (2020).

The climate data were taken from the system of the National Centers for Environmental Forecasting (NCEP), which belongs to the Reanalysis of the Climate Forecasting System (CFRS), where precipitation, wind, relative humidity and solar data were collected in the SWAT file format, from 1979 to 2014. The software used was ArcSWAT, together with ArcGIS 10.3 for the analysis of phosphorus losses.

## Results and discussion

It was observed that crops interfered with the forecast and the way in which nitrogen was lost. Peanut production showed the highest losses according to the simulation, totaling 118.44 kg / ha, while in the sugarcane crop this loss was 34.48 kg / ha. This difference may have occurred because a peanut crop requires more nitrogen than sugarcane, eliminating a greater loss of this nutrient. Legumes are much more demanding, as the root system is superficial and the cycle is specifically shorter, needing to adsorb larger amounts of nitrogen, especially in the no-tillage system when cultivated in succession to grasses [5].

When the total amounts lost from runoff, leaching, lateral flow and nitrate groundwater were compared, the largest losses in sugarcane were observed, 0.46 kg/ha (1.3%), 32.74 kg/ha (95%), 0.19 kg/ha (0.5%) and 0.69 kg/ha (2%), respectively. In the peanut crop, the losses were 0.34 kg/ha (0.3%), 85.99 kg/ha (72.6%), 0.37 kg/ha (0.3%) and 1.84 kg/ha (1.6%), respectively (Table 1).

**Table 01.** Estimate of nitrogen losses in the sub- watershed of Mogi-Guaçu River.

Parameters	Sugarcane (kg/ha)	Percentage (%)	Peanut (kg/ha)	Percentage (%)
<b>Total Nitrogen Loss</b>	34,48	100	118,44	100
<b>Organic nitrogen</b>	0,40	1,20	29,90	25,20
<b>Nitrate superficial runoff</b>	0,46	1,30	0,34	0,30
<b>Leachate nitrate</b>	32,74	95,00	85,99	72,60
<b>Nitrate lateral flow</b>	0,19	0,50	0,37	0,30
<b>Groundwater nitrate</b>	0,69	2,00	1,84	1,60

The main form of nitrogen loss in both production systems occurred through leaching, showing that this process occurred with greater intensity in sugarcane cultivation. Leaching helped in the accumulation of nitrogen in groundwater, being an indicative loss. Nitrate leaching is directly influenced by the factors that determine the flow of water in the soil and the concentration of  $\text{NO}_3^-$  in the solution. In general, the highest concentrations of  $\text{NO}_3^-$  are observed in the most superficial layers of the soil, what is not on the surface usually ends up leaching and accumulating in the groundwater region.  $\text{NH}_4^+$ , on the other hand, are adsorbed by the negative charges in the soil [6].

For the lost amounts of organic nitrogen, the highest were in the peanut crop, totaling 29.90 kg/ha, representing 25.2% of the total loss. As for the sugarcane crop, the amount lost was 0.4 kg/ha, which represented 1.2% of the total loss.

The characteristics of production systems in peanuts promote greater mineralization of organic N due to microbial activity, facilitating the loss of this source, when compared to sugarcane. [7] showed the existence of an increase in N absorption as a function of the greater mineralization of organic N in the soil, the increase in pH, the better distribution of the root

system and also the greater efficiency of symbiotic N<sub>2</sub> fixation due to the lower acidity of the soil. solo [8].

## Conclusions

The Swat tool estimated the nitrogen losses, showing that the management, occupation and landscape of the soil influenced this process, where the biggest losses were in the production of peanuts. Swat can be a management tool, assisting in decisions regarding environmental policies, collaborating with soil and water conservation in Brazilian sub-basins.

**Acknowledgements:** Coordination for the Improvement of Higher Education Personnel (CAPES), Universidade Estadual Paulista (UNESP), Soil Characterization for Specific Management Purposes (CSME) and the Group of Agrometeorological Studies (GAS) for their support.

## References

1. Christoffoleti, P.J., de Carvalho, S.J.P., López-Ovejero, R.F., Nicolai, M., Hidalgo, E., da Silva, J.E. Conservation of natural resources in Brazilian agriculture: implication on weed biology and management. *Crop Protection*, v. 26, p. 383-389. 2007.
2. Brazilian Institute of Geography and Statistics (IBGE). Population Census. 2010. Available at <[www.ibge.gov.br](http://www.ibge.gov.br)>. Accessed on: May 20, 2021.
3. Brazilian Agricultural Research Company (EMBRAPA). National Soil Research Center. Brazilian system of soil classification. Rio de Janeiro, 412 p. 1999.
4. Brazilian Institute of Geography and Statistics (IBGE) - Brazilian Agricultural Research Company (EMBRAPA) - Soil Map of Brazil. Rio de Janeiro: IBGE, Scale 1:5,000,000. 2001. Available at <[www.ibge.gov.br](http://www.ibge.gov.br)>. Accessed on June 27, 2020.
5. Gomes Júnior, F.G.; Sa, M.E.; Valério, W.V. Nitrogen in common bean under no-tillage system on grasses. *Acta Scientiarum Agronomy, Maringá*, v. 30, no. 3, p. 387-395, 2008.
6. Lai, X.; Zhou, Z.; Liao, K.; Zhu, Q. Responses of soil carbon and nitrogen cycles to the physical influences of rock fragment in soils. *CATENA*, v. 203, p. 105369, 2021.
7. Rosolem, C.A.; Pereira, H.F.M.; Bessa, A.M.; Amaral, P.G. Nitrogen in soil and cotton growth as affected by liming and N fertilizer. In: Wright, R.J.; Baligar, V.C.; Murrmann, R.P. (Ed.) *Plant-soil interactions at low pH*. Dordrecht: Kluwer, p.321-325, 1990.
8. Blamey, F.P.C. & Chapman, J. Soil amelioration effects on peanut growth, yield and quality. *Plant and Soil*, The Hague, vol. 65, p. 319-334, 1982.